



# Measurement of Heavy Metals (Nickel, Chromium, and Cobalt) in Wild and Farmed Carps (*Cyprinus carpio*) of Hamadan Province

Mohammad Moghadasi<sup>1</sup>, Ali Heshmati<sup>2</sup>, Aliasghar Vahidinia<sup>3\*</sup>

<sup>1</sup>Student Research Center, Hamadan University of Medical Sciences, Hamadan, Iran

<sup>2</sup>Department of Nutrition and Food Hygiene, School of Medicine, Nutrition Health Research Center, Hamadan University of Medical Sciences, Hamadan, Iran

<sup>3</sup>Department of Nutrition and Food Hygiene, Nutritional Health Research Center, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

## \*Correspondence to

Aliasghar Vahidinia,  
Email: vahidinia@umsha.ac.ir

Published online December 29, 2021

## Abstract

Fish consumption has been considerably increased in Iran recently. On the other hand, the increase in aquatic ecosystem pollution can cause the accumulation of heavy metals in aquatic animals – the fish, in particular. Therefore, measuring the amount of heavy metals in fish is of great importance for consumers' health. This study aimed to measure the amount of heavy metals (nickel, chromium, and cobalt) in wild and farmed carps (*Cyprinus carpio*) by using an ICP device. In this cross-sectional study which was performed in Hamadan province in 2018, 40 carp samples (20 wild carp samples and 20 farmed ones) were randomly collected from Anouch and Shirinsoo dams and from breeding centers of the province and, after completing the digestion phase, the amount of heavy metals in the samples were measured using the ICP device. The study results showed that the amounts of nickel metal in wild and farmed carps were equal to  $0.108 \pm 0.086$  and  $0.102 \pm 0.108$  mg/kg dry weight, respectively; and the amounts of chromium in wild and farmed fish samples were  $0.348 \pm 0.396$  and  $0.136 \pm 0.074$  mg/kg dry weight in muscle tissue, respectively. Cobalt metal was not observed in wild and cultured samples. According to the results from the statistical analysis, no significant difference was detected between wild and farmed fishes regarding contamination with nickel metal. However, a difference was found between wild and farmed fishes ( $P < 0.05$ ) regarding chromium. Taking into account the higher amount of nickel and chromium metals in the wild carp samples compared to the farmed ones, it was recommended that necessary measures be taken to control the sources responsible for polluting the aquatic environment and to prevent the penetration of industrial effluents and other polluting factors into the dams of Hamedan province in order for avoiding the increase in the amount of heavy metals in the province's water sources.

**Keywords:** Cobalt, Chrome, Nickel, *Cyprinus carpio*, Heavy metals

Received July 19, 2021; Revised September 29, 2021; Accepted October 20, 2021



## 1. Introduction

Heavy metal toxicity poses a significant threat to the environment and public health since it increases the pollution level which, in turn, causes serious problems for living organisms. The second sentences should be replaced with the following phrase: Therefore, the information about their level in various products consumed living organism is important (1, 2). In health sciences, fish is argued to serve as a reliable indicator for assessing heavy metal pollution level in seas, dams, and rivers (3). These metals are one of the most important pollutants that harm aquatic environment and seriously endanger the health of aquatic living beings by, for example, poisoning the fish (4). Aquatic environment pollution is usually caused by the industrial effluents, sewage, petroleum products,

and gasoline from ships and boats (5). Nowadays, aquatic products – fish, in particular, are considered as rich protein sources for human body. In addition, fish provides human body with rich sources of vitamin and omega-3 fatty acids (6). Common carp is one of the most commercially important fish in the world which is farmed commercially and wildly. Carps are omnivorous, but usually feed on sediments in the water and may accumulate more heavy metals than other types of fish (7). Farmed fishes have been noted for their freshness and low price. Eating fish twice a week has been highly recommended by the American Heart Association (8). Heavy metals in effluents from factories and industrial sites may find their way into seawater and pollute the aquatic ecosystem. These metals are resistant to decomposition

and can accumulate in the aquatic living organisms (9). Chromium, cobalt, and nickel are among potentially toxic elements (PTEs) found in the aquatic environment, and the World Organization has warned against exposure to them (10). Chromium is a toxic and unnecessary metal found in soil and water sources, which can pose a serious threat to human health through the food chain. One of the side effects of multiple exposure to chromium (VI) is liver toxicity in human (11, 12). Chromium exists in two major stable oxidation states, hexavalent chromium (Cr (VI)) and trivalent chromium (Cr (III)) (13, 14). Cobalt is one of the most dangerous metals for living organisms, which is stable in the forms of  $Co + 3$  and  $Co + 2$  (15). Cobalt constituents are believed to cause cancer and respiratory tumors in human (16). Nickel metal which is a toxic element found in aquatic environment, air, and soil can cause respiratory cancer, dermatitis, and reproductive problems for human (2). Younesipour et al found that the levels of nickel and cobalt in the Caspian Sea carps were equal to  $121 \pm 14.3$  and  $323.5 \pm 28.34$  ( $\mu\text{g/g}$  dry weight), respectively (17). In the study by Solgi et al and Sarvai et al measuring the levels of heavy metals in common carp, it was revealed that the amounts of cobalt and nickel were lower than the limits defined by WHO (18-20). Heshmati et al also attempted to examine the levels of heavy metals in the Caspian Sea and discovered that the levels of nickel and cobalt in wild and farmed carps were not significantly different (21). In a study by Varol et al in Turkey, it was determined that the concentrations of cobalt, nickel, and chromium in carps were equal to  $0.081 \pm 0.035$ ,  $0.234 \pm 0.407$ , and  $0.074 \pm 0.074$  (mg/kg), respectively (22). According to that, little information was available about the amount of heavy metals in wild and farmed carp species in Hamadan province. Moreover, the evidence showed that some pesticides and environmental pollutants had penetrated into the Shirinsoo and Anouch dams due to the increasing agricultural activities and their water had been exposed to rural and urban sewage, posing a serious threat to carp consumers in the province due to the increasing toxicity of farmed fish. This study, therefore, aimed to measure the amount of heavy metals nickel, chromium, and cobalt in wild and farmed carps.


## 2. Materials and Methods

### 2.1. Study Area and Time Description

The areas investigated in the present study were Shirinsoo and Anouch dams and carp breeding centers of Hamadan province in the autumn of 2018. The geographical locations of the fishing sites are shown in Fig. 1.

### 2.2. Sample Collection

A total of 40 carp samples (20 wild carp samples from Shirinsoo and Anouch dams, and 20 farmed carp samples from fish breeding centers) were randomly collected and



Wild carp caught from Shirin Su Dam ↑

Farmed carp caught from Anouch Dam ↑

**Fig. 1.** Geographical Location of Carp Samples Caught in Shirinsoo and Anouch Dams of Hamadan Province. White arrow = Wild Carp Caught From Shirin-Su Dam; Black arrow = Farmed Carp Caught From Anouch Dam.

transferred to the Food and Nutrition Control Laboratory of Hamadan University of Medical Sciences and, then, they were stored in a freezer at minus  $18^{\circ}\text{C}$ .

### 2.3. Sample Preparation

Materials used in this study included hydrochloric acid 37%, nitric acid 65%, and hydrogen peroxide, all of which were purchased from the Merck Co., Germany. Prior to the experiment, the frozen fish were thawed and their weight and length were determined by a digital scale and a simple ruler in the next step. The means and standard deviations of length and weight of wild and farmed carps are shown in Table 1. Then some of the fish's edible muscles were removed, weighed, and placed on a clean plate. The plate containing the sample was transferred to an oven at  $115^{\circ}\text{C}$  and kept in it for 120 minutes to let the sample dry completely. The dried fish weight was measured and its moisture content was determined (21). The means and standard deviations of the dry and wet matter of wild and farmed carps are shown in Table 2. To complete the digestion phase, 1 g of the dried fish muscle was placed in Erlenmeyer and 5 cc of a mixture of 37% hydrochloric acid and 65% nitric acid in a ratio of 3 to 1 was added to Erlenmeyer; then it was placed on the heater at  $95^{\circ}\text{C}$  for 60 minutes. After 60 minutes, a mixture of 5 cc of hydrochloric acid and nitric acid was added to Erlenmeyer again and the heating by the heater was continued for another hour until a clear color appeared. Then 2 cc of hydrogen peroxide was added to the Erlenmeyer to make the color of the Erlenmeyer contents completely clear. After cooling Erlenmeyer, the samples were filtered using filter paper and bring up to the 25 mL in a Falcon tube. Heavy metals were measured by the ICP-OES (inductively coupled plasma - optical emission spectrometry) method using the ARCOS FHE12 model

**Table 1.** Means and Standard Deviations of Length (cm) and Weight (g) of Wild and Farmed Carps Studied in Hamadan Province in 2018

Crap ( <i>Cyprinus carpio</i> )	Weight (g) Mean $\pm$ SD	The Length Mean $\pm$ SD
Wild crap	76.480 $\pm$ 52.305	18.05 $\pm$ 3.48
Farmed crap	154.87 $\pm$ 127.894	22.15 $\pm$ 6.607
P value	0.006	0.028

**Table 2.** Means and Standard Deviations of Moisture Content (%) and Dry Matter (%) in Wild and Farmed Fish Studied in Hamadan Province in 2018

Crap ( <i>Cyprinus carpio</i> )	Dry matter (%) Mean $\pm$ SD	Moisture (%) Mean $\pm$ SD
Wild crap	76.542 $\pm$ 4.694	23.457 $\pm$ 4.577
Farmed crap	75.537 $\pm$ 5.557	24.463 $\pm$ 5.557
P value	0.07	0.213

(AMETEK company, Germany) (23).

#### 2.4. Statistical Analysis

SPSS software version 16 was employed for statistical analysis and Kolmogorov-Smirnov test was performed for determining the normal distribution of data. Man-Whitney test was also used to compare the data without normal distribution. If ( $P < 0.05$ ), a significant difference was assumed.

### 3. Results and Discussion

The linearity range and calibration curve equation for the three metals are shown in Table 3. The *limit of detection* (LODs) of Ni, Cr, and Co were 0.000454, 0.000471, and 0.00208 (mg/kg<sup>-1</sup>) respectively. The average concentrations of nickel in wild and farmed carps were equal to 0.086  $\pm$  0.108 and 0.108  $\pm$  0.102 mg/kg dry weight, respectively; and the average concentrations of chromium in muscle tissues of wild and farmed samples were equal to 0.396  $\pm$  0.348 and 0.074  $\pm$  0.136 mg/kg dry weight, respectively. However, cobalt metal was not detected in free and cultured carp samples, and its level was below the detection limit of the device. In this study, the amounts of metals measured in salmon and farmed fish in the forms of chromium > nickel > cobalt were also obtained. The amounts of chromium and nickel metals in wild carp samples were higher than the amount found in farmed carp samples; and the amounts of the given metals in wild carp samples were higher than those detected in farmed carp samples, which were lower than the global

standards determined by of FAO, WHO, and FDA (24-26). Also, the amount of chromium metal in both farmed and wild carps was lower than the limit defined by global FDA (26). According to our study results from statistical analysis, no significant difference was observed between the edible tissues of wild and farmed fishes in terms of the concentration of nickel metal, while a noticeable difference was observed ( $P < 0.05$ ) between wild and farmed fishes regarding chromium metal concentration. This significant difference may have been due to the presence of surface sediments and/or the entry of rural/agricultural effluents and toxic substances such as pesticides from adjacent agricultural lands into dams water (27,28). The mean and standard deviations of the wild and farmed carps are shown in Table 4. Due to the special importance attached to the determination of heavy metals in various food products in Iran and other parts of the world – wild and farmed fishes, in particular, this study aimed to investigate the concentrations of nickel, chromium, and cobalt in samples from wild and farmed carps. Chromium is one of the toxic metals which can be accumulated and disrupted in the host body. Nickel and cobalt are essential metals for the body and their excessive intake is harmful to human organs (29). In the studies by Nasrollahzadeh Saravi et al and Mansouri et al, the amounts of nickel metal and cobalt in the muscle tissues of carps had been discovered to be very insignificant and lower than the limits determined by FAO international standards, which were consistent with our study findings (19,30). Our study findings showed that the amount of nickel metals in wild carp was higher than that detected in farmed carp, but no significant difference was observed. This finding was in line with the finding from the study by Heshmati et al (21). Nickel is one of the most abundant elements in the ecosystem, whose high levels in wild fish can be due to the presence of silicate minerals and water pollution caused by industrial wastewater (31). In a

**Table 4.** Means and Standard Deviations (mg/kg dry weight) of the Studied Metals in Muscle Tissue of Wild and Farmed Fish in Hamadan Province in 2018

Metal	Farmed Crap (mg/kg)	Wild Crap (mg/kg)	P value
Nickel	0.108 $\pm$ 0.102	0.108 $\pm$ 0.086	0.54
Chromium	0.074 $\pm$ 0.136	0.348 $\pm$ 0.396*	0.002
Cobalt	ND	ND	ND

\* The results have significant differences.  
ND: not detected.

**Table 3.** Linear Equations and Regression Coefficient of the Calibration Curves and LOD and LOQ of Studied and Semi-essential Metals

LOQ (mg/kg)	LOD (mg/kg)	Regression Coefficient	Equation	Range (mg/L)	Metals
0.00154	0.000454	0.99999	Y=0.00002.996X-0.2065	0.000454-6	Ni
0.00160	0.000471	1	Y=0.000005.4454X-0.0744	0.000471-6	Cr
0.00707	0.00208	1	Y=0.000006.551X-0.0944	0.00208-6	Co

LOD, Limit of detection; LOQ, limit of quantification.

study by Kalyoncu et al, the amount of chromium in carp samples was found to have been higher than that of the nickel in the given fish, which was consistent with result from the present study (32). Chromium metal is known as a toxic element, and its increase in the environment and aquatic ecosystem may be due to the widespread use of pesticides near the seas, rivers, and dams (33, 34). In the study by Leung et al, however, the amount of nickel metal in the examined fish was detected to have been higher than that of chromium metal, which was not consistent with our study findings in this regard (35). The high amount of nickel metal found by this study may have been due to pollution caused by oil as well as the traffic of oil ships and motorboats Pearl river delta/china (36). In Elsagh's study, the concentration of chromium metal in carp from the Caspian Sea was reported to be higher than the level determined by international standards, which was contrary to the result obtained in the present study; this inconsistency may have been due to the higher pollution of the Caspian Sea and enormous increase in using pesticides in this area compared to water basins of Hamadan province (37). In a study by Tkachenko et al, the amount of chromium metal was determined to be higher than that of cobalt metal in farmed carp in Poland, which was in agreement with our study results (38). Salemi and Hosseini Alhashemi reported that the concentration of chromium in common carp from the shores of Dez was equal to 0.02 mg/kg dry weight and was lower than the limit defined by international standards, which was consistent with our study findings; however, they demonstrated that the concentration of nickel in this fish was equal to 1.41 mg/kg dry weight and was higher than the limit defined by international standards, which was inconsistent with the findings from this study (39). In another study by Saleem et al, it was found that the concentration of cobalt metal in Pakistani carp samples was higher than the amount of nickel and chromium, which was contrary to the results from our study since cobalt metal was not observed in wild and farmed carp samples examined in our study and its amount was below the detection limit of the device (40). The high amount of cobalt recorded by Saleem et al may have been attributed to the existence of agricultural lands and the extensive use of fertilizers and chemical materials in the investigated area (41). In a study by Ahmadi, no concentration of cobalt metal was observed in the *Esox lucius* from west of Anzali wetland and it was not within the detection of the device, which was consistent with our study findings (42).

#### 4. Conclusion

Although the amount of cobalt metal was below the detection limit and the chromium level in the tested samples was lower than the one defined by the world standards, there was a possibility that the excessive use of pesticides as well as the agricultural discharge of

effluents to nearby lands, dams, and water sources in fish farms would result in an increase in the amount of heavy metals in the given areas in the following years. Therefore, it was suggested that the local people and farmers be educated about the dangers of heavy metals, and the entry of environmental pollutants into the dams of Hamadan province be effectively controlled as much as possible. As the amounts of nickel and chromium metals in wild carp samples were higher than those in farmed carp samples, moreover, it was recommended that the necessary measures be taken in order to control the factors responsible for polluting the aquatic environment, and to prevent the entry of industrial effluents and other pollutants into the dams of Hamadan province for avoiding the increase in the amount of heavy metals in water sources.

#### Conflict of Interest Disclosures

The authors declare that they have no conflict of interests.

#### Acknowledgment

Authors would like to thank all those people who contributed to this research, specially Mrs. Janaati for helping us in different stages of this research work. This study was funded by the Hamadan University of Medical Sciences, Hamadan, Iran (No. 9711026527).

#### References

1. Saeedi Saravi SS, Karami S, Karami B, Shokrzadeh M. Toxic effects of cobalt chloride on hematological factors of common carp (*Cyprinus carpio*). Biol Trace Elem Res. 2009;132(1-3):144-52. doi: [10.1007/s12011-009-8388-8](https://doi.org/10.1007/s12011-009-8388-8).
2. Sachan P, Lal N. An overview of nickel (Ni<sup>2+</sup>) essentiality, toxicity and tolerance strategies in plants. Asian J Biol. 2017;2(4):1-15. doi: [10.9734/ajob/2017/33931](https://doi.org/10.9734/ajob/2017/33931).
3. Kalani N, Karbassi AR, Moatar F. Evaluation of Heavy Metals Arsenic Lead Cadmium Chromium and Nickel in Muscle Liza saliens and Evaluation of Health Risk Due to Its Consumption. Journal of Aquaculture and Fisheries. 2014;5(17):65-79.
4. Authman MM, Zaki MS, Khallaf EA, Abbas HH. Use of fish as bio-indicator of the effects of heavy metals pollution. J Aquac Res Dev. 2015;6(4):328. doi: [10.4172/2155-9546.1000328](https://doi.org/10.4172/2155-9546.1000328).
5. Velusamy A, Satheesh Kumar P, Ram A, Chinnadurai S. Bioaccumulation of heavy metals in commercially important marine fishes from Mumbai Harbor, India. Mar Pollut Bull. 2014;81(1):218-24. doi: [10.1016/j.marpolbul.2014.01.049](https://doi.org/10.1016/j.marpolbul.2014.01.049).
6. Adeli A. Properties of fish and its nutritional value for humans. Journal of Fisheries, Islamic Azad University, Azadshahr Branch. 2013;9(3):61-8.
7. Sobhanardakani S, Jafari SM. Investigation of As, Hg, Zn, Pb, Cd and Cu concentrations in muscle tissue of *Cyprinus carpio*. J Mazandaran Univ Med Sci. 2014;24(116):184-95. [Persian].
8. Özogul Y, Özogul F, Gökbulut C. Quality assessment of wild European eel (*Anguilla anguilla*) stored in ice. Food Chem. 2006;95(3):458-65. doi: [10.1016/j.foodchem.2005.01.025](https://doi.org/10.1016/j.foodchem.2005.01.025).
9. Najm M, Shokrzadeh M, Fakhar M, Sharif M, Hosseini SM, RahimiEsboei B, et al. Concentration of heavy metals (Cd, Cr and Pb) in the tissues of *Clupeonella cultriventris* and *Gasterosteus aculeatus* from Babolsar coastal waters of Mazandaran province, Caspian Sea. J Mazandaran Univ Med Sci. 2014;24(113):185-92. [Persian].
10. Ustaoglu F, Islam MS. Potential toxic elements in sediment of some rivers at Giresun, Northeast Turkey: a preliminary assessment for ecotoxicological status and health risk. Ecol Indic. 2020;113:106237. doi: [10.1016/j.ecolind.2020.106237](https://doi.org/10.1016/j.ecolind.2020.106237).

11. Shahid M, Shamshad S, Rafiq M, Khalid S, Bibi I, Niazi NK, et al. Chromium speciation, bioavailability, uptake, toxicity and detoxification in soil-plant system: a review. *Chemosphere*. 2017;178:513-33. doi: [10.1016/j.chemosphere.2017.03.074](https://doi.org/10.1016/j.chemosphere.2017.03.074).
12. Tian X, Zhang H, Zhao Y, Mehmood K, Wu X, Chang Z, et al. Transcriptome analysis reveals the molecular mechanism of hepatic metabolism disorder caused by chromium poisoning in chickens. *Environ Sci Pollut Res Int*. 2018;25(16):15411-21. doi: [10.1007/s11356-018-1653-7](https://doi.org/10.1007/s11356-018-1653-7).
13. Sarkar S, Satheshkumar A, Premkumar R. Hexavalent chromium (Cr(VI)) removal by live mycelium of a *Trichoderma harzianum* strain. *Mol Soil Biol*. 2013;4(1):1-6. doi: [10.5376/msb.2013.04.0001](https://doi.org/10.5376/msb.2013.04.0001).
14. Fang Z, Zhao M, Zhen H, Chen L, Shi P, Huang Z. Genotoxicity of tri- and hexavalent chromium compounds in vivo and their modes of action on DNA damage in vitro. *PLoS One*. 2014;9(8):e103194. doi: [10.1371/journal.pone.0103194](https://doi.org/10.1371/journal.pone.0103194).
15. Javed M, Usmani N. An overview of the adverse effects of heavy metal contamination on fish health. *Proc Natl Acad Sci India Sect B Biol Sci*. 2019;89(2):389-403. doi: [10.1007/s40011-017-0875-7](https://doi.org/10.1007/s40011-017-0875-7).
16. Lison D, van den Brule S, Van Maele-Fabry G. Cobalt and its compounds: update on genotoxic and carcinogenic activities. *Crit Rev Toxicol*. 2018;48(7):522-39. doi: [10.1080/10408444.2018.1491023](https://doi.org/10.1080/10408444.2018.1491023).
17. Unesipour H, Nasrollahzadehsaravi H, Saddatipour MT. Investigation of bioaccumulation of essential heavy metals iron, copper and zinc (and semi-essential) nickel, cobalt and manganese in the edible tissue of carp (*Cyprinus carpio*) Caspian Sea. *Journal of Aquaculture Development*. 2013;8(1):95-106.
18. Solgi E, Alipour H, Majnoon F. Evaluation of heavy metal concentrations in common carp muscle on the southern shores of the Caspian Sea and the potential risk to human health. *Iranian Journal of Fisheries*. 2017;27(1):113-123.
19. Nasrollahzadeh Saravi H, Pourgholam R, Pourang N, Rezaei M, Makhloogh A, Unesipour H. Heavy metal concentrations in edible tissue of *Cyprinus carpio* and its target hazard quotients in the Southern Iranian Caspian Sea coast, (2010). *J Mazandaran Univ Med Sci*. 2013;23(103):33-44. [Persian].
20. World Health Organization (WHO). Guidelines for Drinking-Water Quality. WHO; 1993.
21. Heshmati A, Karami-Momtaz J, Nili-Ahmadabadi A, Ghadimi S. Dietary exposure to toxic and essential trace elements by consumption of wild and farmed carp (*Cyprinus carpio*) and Caspian kutum (*Rutilus frisii kutum*) in Iran. *Chemosphere*. 2017;173:207-15. doi: [10.1016/j.chemosphere.2017.01.009](https://doi.org/10.1016/j.chemosphere.2017.01.009).
22. Varol M, Kaya GK, Sünbül MR. Evaluation of health risks from exposure to arsenic and heavy metals through consumption of ten fish species. *Environ Sci Pollut Res Int*. 2019;26(32):33311-20. doi: [10.1007/s11356-019-06450-x](https://doi.org/10.1007/s11356-019-06450-x).
23. American Public Health Association (APHA), American Water Works Association (AWWA), Water Pollution Control Federation, Water Environment Federation (WEF). Standard Methods for the Examination of Water and Wastewater. APHA; 1912.
24. World Health Organization (WHO). Obesity: Preventing and Managing the Global Epidemic. WHO; 2000.
25. Nauen CE. Compilation of Legal Limits for Hazardous Substances in Fish and Fishery Products. Rome: FAO; 1983.
26. Food and Drug Administration (FDA). Fish and Fisheries Products Hazards and Controls Guidance. Washington, DC: Food and Drug Administration, Center for Food Safety and Applied Nutrition; 2001.
27. Gagnon C, Fisher NS. The bioavailability of sediment-bound Cd, Co, and Ag to the mussel *Mytilus edulis*. *Can J Fish Aquat Sci*. 1997;54(1):147-56. doi: [10.1139/f96-256](https://doi.org/10.1139/f96-256).
28. Morillo J, Usero J, Gracia I. Heavy metal distribution in marine sediments from the southwest coast of Spain. *Chemosphere*. 2004;55(3):431-42. doi: [10.1016/j.chemosphere.2003.10.047](https://doi.org/10.1016/j.chemosphere.2003.10.047).
29. Norouzi M, Bagheri Tavani M. Assessment of the consumption risk of fifteen heavy metals in *Liza aurata* in the Caspian Sea. *J Aquat Ecol*. 2019;8(3):96-107. [Persian].
30. Mansouri B, Majnoni F, Maleki A, Rezaei Z, Azadi N, Gharibi F. Bioaccumulation of Pb, Cd, Zn, and Cu in the muscle, gill, liver, and skin of common carp (*C. carpio*) and silver carp (*H. molitrix*) in Geshlagh dam in Sanandaj city. 2014. *Zanko J Med Sci*. 2014;15(45):26-35. [Persian].
31. Naji T, Safaeian SH, Rostami M, Sabrjou M. Toxic effect of zinc sulfate on gill tissues of common carp (*Cyprinus carpio*). *J Environ Sci Technol*. 2007;9(233):29-36. [Persian].
32. Kalyoncu L, Kalyoncu H, Arslan G. Determination of heavy metals and metals levels in five fish species from Işıklı Dam Lake and Karacaören Dam Lake (Turkey). *Environ Monit Assess*. 2012;184(4):2231-5. doi: [10.1007/s10661-011-2112-9](https://doi.org/10.1007/s10661-011-2112-9).
33. Michalak I, Zielińska A, Chojnacka K, Matuła J. Biosorption of Cr(III) by microalgae and macroalgae: equilibrium of the process. *Am J Agric Biol Sci*. 2007;2(4):284-90.
34. Sabbagh A. Determine the Level of Heavy Metals in Muscle, Liver, Gills, Kidney and Ovarian *Lizza aurata* on the Southern Coast of the Caspian Sea [thesis]. Tehran: Environmental Sciences, University of Tarbiat Modares; 2001. [Persian].
35. Leung HM, Leung AO, Wang HS, Ma KK, Liang Y, Ho KC, et al. Assessment of heavy metals/metalloid (As, Pb, Cd, Ni, Zn, Cr, Cu, Mn) concentrations in edible fish species tissue in the Pearl River Delta (PRD), China. *Mar Pollut Bull*. 2014;78(1-2):235-45. doi: [10.1016/j.marpolbul.2013.10.028](https://doi.org/10.1016/j.marpolbul.2013.10.028).
36. Tabari S, Saeedi Saravi SS, Bandany GA, Dehghan A, Shokrzadeh M. Heavy metals (Zn, Pb, Cd and Cr) in fish, water and sediments sampled from Southern Caspian Sea, Iran. *Toxicol Ind Health*. 2010;26(10):649-56. doi: [10.1177/0748233710377777](https://doi.org/10.1177/0748233710377777).
37. Elsagh A. Bioaccumulation of heavy metals levels in muscles of *Rutilus frisii kutum* and *Cyprinus carpio* fishes of coastal waters of the Mazandaran province, Caspian Sea. *Veterinary Researches & Biological Products*. 2012;25(2):41-8. doi: [10.22092/vj.2012.101069](https://doi.org/10.22092/vj.2012.101069). [Persian].
38. Tkachenko H, Kurhaluk N, Kasiyan O, Kamiński P. Dietary nutrients and health risks from exposure to some heavy metals through the consumption of the farmed common carp (*Cyprinus carpio*). *J Environ Health Sci Eng*. 2021;19(1):793-804. doi: [10.1007/s40201-021-00647-4](https://doi.org/10.1007/s40201-021-00647-4).
39. Salemi M, Hosseini Alhashemi AA. Bioaccumulation of heavy metals (cadmium, chromium, nickel, zinc), in *Cyprinus carpio* fish. *J Anim Biol*. 2017;9(4):35-45.
40. Saleem M, Shi Z, Shah MH. Evaluation of contamination status and health risk assessment of essential and toxic metals in *Cyprinus carpio* from Mangla Lake, Pakistan. *Biol Trace Elem Res*. 2021;199(11):4284-94. doi: [10.1007/s12011-020-02540-x](https://doi.org/10.1007/s12011-020-02540-x).
41. Lado LR, Hengl T, Reuter HI. Heavy metals in European soils: a geostatistical analysis of the FOREGS geochemical database. *Geoderma*. 2008;148(2):189-99. doi: [10.1016/j.geoderma.2008.09.020](https://doi.org/10.1016/j.geoderma.2008.09.020).
42. Ahmadi M. Evaluation of copper and cobalt heavy metals in edible duck muscle tissue of Bandar Anzali International Wetland. *Aquatic Exploitation and Breeding*. 2020;9(2):13-20.